

Invasion of temperate riparian forests by *Acacia dealbata* affects macroinvertebrate community structure in streams.

Silvia Gómez-Arcusa^{1*} , Nuno Coimbra² and Verónica Ferreira² 

¹ Department of Life Sciences, University of Coimbra, Calçada Martim de Freitas, 3000–456 Coimbra, Portugal.

² MARE – Marine and Environmental Sciences Centre, ARNET – Aquatic Research Network, Department of Life Sciences, University of Coimbra, Calçada Martim de Freitas, 3000–456 Coimbra, Portugal.

* Corresponding author: sgomezarcusa@gmail.com

Received: 4/09/23

Accepted: 11/01/24

ABSTRACT

Invasion of temperate riparian forests by *Acacia dealbata* affects macroinvertebrate community structure in streams.

The riparian forest determines many of the stream characteristics such as physicochemical variables and the availability and quality of food resources for aquatic communities. In central Portugal, native mixed deciduous forests are being heavily invaded by *Acacia dealbata*, which is an evergreen and nitrogen-fixing tree species, but effects on stream benthic macroinvertebrate communities are unknown. We assessed invasion effects on macroinvertebrate communities by comparing streams flowing through native forests (native streams) and streams flowing through forests heavily invaded by *A. dealbata* (invaded streams), in central Portugal. We did not find major differences in macroinvertebrate abundance, total taxa richness or diversity between native and invaded streams, due to high variation among native streams. However, taxonomic structure of macroinvertebrate communities differed between stream types, with invaded streams showing lower percentage of sensitive taxa. Macroinvertebrate communities structure based on functional feeding groups also differed between stream types, with a tendency for higher abundance of shredders and scrapers/grazers and lower abundance of filter feeders in native than invaded streams, probably due to altered composition of leaf litter inputs and canopy cover within invaded streams. Our findings suggest that invasion by *A. dealbata* affects the macroinvertebrate community mostly in terms of taxonomic and functional structure. Since *A. dealbata* is rapidly spreading in all southern Europe, this study highlights the importance of enhancing the comprehension of its impact on aquatic communities to identify ecological risks and formulate effective mitigation strategies.

KEYWORDS: Alien vegetation, benthic macroinvertebrates, FFG, forest change, functional structure, litter supply, N-fixing tree

RESUMEN

La invasión de bosques ribereños templados por *Acacia dealbata* afecta la estructura de la comunidad de macroinvertebrados en arroyos.

El bosque ribereño determina muchas de las características de los arroyos, como las variables fisicoquímicas y la disponibilidad y calidad de los recursos para las comunidades acuáticas. En el centro de Portugal, los bosques mixtos autóctonos de hoja caduca están siendo fuertemente invadidos por *Acacia dealbata*, un árbol perenne fijador de nitrógeno, pero se desconocen los efectos sobre las comunidades de macroinvertebrados bentónicos de los arroyos. Hemos evaluado los efectos de la invasión sobre las comunidades de macroinvertebrados mediante la comparación entre arroyos que fluyen a través de bosques nativos (arroyos nativos) y arroyos que fluyen a través de bosques invadidos por *A. dealbata* (arroyos invadidos), en el centro de Portugal. No se han encontrado diferencias significativas en la abundancia, riqueza total de taxones y diversidad de macroinvertebrados entre arroyos nativos e invadidos, debido a la alta variación entre arroyos nativos. Sin embargo, la estructura taxonómica de las comunidades de macroinvertebrados difirió entre los dos tipos

de arroyos, con un menor porcentaje de taxones sensibles en los arroyos invadidos. La estructura de las comunidades basada en grupos de estrategia trófica también difirió entre los tipos de arroyos, con tendencia a una mayor abundancia de trituradores y raspadores/pastadores y una menor abundancia de filtradores en los arroyos nativos en comparación con los arroyos invadidos, probablemente debido a la alteración de los inputs de hojarasca y de la cobertura de dosel arbóreo en los arroyos invadidos. Nuestros hallazgos sugieren que la invasión por *A. dealbata* afecta principalmente a la comunidad de macroinvertebrados en términos de estructura taxonómica y funcional. Dado que *A. dealbata* se está propagando rápidamente en todo el sur de Europa, este estudio resalta la importancia de mejorar la comprensión de su impacto en las comunidades acuáticas para identificar riesgos ecológicos y formular estrategias de mitigación.

PALABRAS CLAVES: árbol fijador de nitrógeno, aporte de hojarasca, cambio de bosque, estructura funcional, FFG, macroinvertebrados bentónicos, vegetación exótica.

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.

INTRODUCTION

Streams and their riparian forests are closely connected ecosystems, linked by the bidirectional flow of matter and energy; e.g., plant litter flows from the riparian forest into the stream where it sustains the detrital food web, and adult aquatic insects flow in the opposite direction to integrate terrestrial food webs (Tolkkinen *et al.*, 2020). The riparian forest determines many of the stream characteristics; e.g., canopy cover influences solar exposure of the stream bed and water temperature (Beschta, 1997), nutrient uptake by and litter inputs from the riparian forest affect the water chemical composition (Dosskey *et al.*, 2010) and the latter also affects the benthic organic matter standing stock (Abelho & Graça, 1996; Molinero & Pozo, 2004), and the supply of branches and logs to streams affects current velocity and depth conditions and changes microhabitat distribution (Gerhard & Reich, 2000). Consequently, the riparian forest influences the instream energetic pathways (autotrophic vs. heterotrophic) and the structure of aquatic communities (Hawkins *et al.*, 1982; Ferreira *et al.*, 2016; Little *et al.*, 2021).

The invasion of native riparian forests by exotic tree species strongly affects forest tree composition and, consequently, the amount, diversity, taxonomic and chemical composition, and seasonality of the litter supply to streams (Lecerf *et al.*, 2007; Hladyz *et al.*, 2011; Mineau *et al.*, 2012). Riparian forest invasion also affects stream characteristics such as water nutrient concentrations; e.g., increase in the cover by nitrogen-fixing species generally leads to increases in dissolved nitrogen concentration in stream wa-

ter (Goldstein *et al.*, 2009; Atwood *et al.*, 2010; Stewart *et al.*, 2019). Changes in food resources (i.e., litter inputs) and stream water characteristics resulting from forest invasion can potentially affect aquatic communities. For example, the invasion of the Fynbos vegetation by *Acacia mearnsii* De Wild. in South Africa resulted in lower taxa richness (Magoba & Samways, 2010) and altered benthic macroinvertebrate community structure, with lower abundance of cobble-dwelling taxa and higher abundance of particle-feeding taxa in stream sites flowing through invaded than through native vegetation (Lowe *et al.*, 2008). Invasion of temperate deciduous forests by *Lonicera maackii* (Rupr.) Maxim. in the American Midwest also led to changes in benthic macroinvertebrate community structure along an invasion gradient, with the most invaded stream showing the lowest percentage of sensitive taxa (Little *et al.*, 2021). In central and southern Scotland, invasion of riparian areas by exotic plants was associated with a decrease in the diversity and an increase in the abundance of stream benthic macroinvertebrates, and changes in community structure in autumn (but not in spring) (Seeney *et al.*, 2019). Finally, invasion of riparian areas by *Salix fragilis* L. in the northern Patagonian Andes led to changes in stream benthic macroinvertebrate community structure but not in abundance and diversity (Serra *et al.*, 2013). These examples, although not intended to be extensive, show that invasion effects somewhat vary among studies that differ in the native vegetation, invasive species, climate, and potentially also in the extent and duration of the invasion. Still, effects of riparian invasions were pervasive for stream benthic macroinvertebrate

community structure. Also, effects of species invasions are likely stronger when native and invasive species show different functional characteristics (Kominoski et al., 2013; Castro-Díez et al., 2014).

In central Portugal, native forests dominated by deciduous tree species (*Quercus* spp., *Castanea sativa* Mill.) are being heavily invaded by *Acacia* species originating from Australia, mainly *A. dealbata* Link.; presently, *Acacia* species cover up to 43% of the area in some small-to-medium sized stream basins (Ferreira et al., 2021; Pereira et al., 2021) (Fig. 1). The invasion success of *Acacia* species is due to its (i) great plasticity that allows colonization of disturbed soils (e.g., stream banks) (Pohlman et al., 2005), (ii) ability to fix atmospheric nitrogen (N) that allows high growth rate in nutrient-poor and acidic soils (Gallagher et al., 2011), (iii) high seed production that confers high spread ability (Lorenzo et al., 2010), and (iv) production of allelopathic compounds that inhibit the germination and growth of competitor species (Lorenzo et al., 2008). In addition to covering large areas (Fig. 1), *A. dealbata* is an evergreen and N-fixing species, traits that markedly differ from those of the dominant native tree species (Lorenzo et al., 2010; Ferreira et al., 2021). Therefore, invasion of native diverse deciduous forests with low representation of N-fixing tree species by *A. dealbata* can potentially

affect aquatic communities of adjacent streams (Ferreira et al., 2021).

In this study we compared the benthic macroinvertebrate communities between streams flowing through native mixed deciduous forests and streams flowing through forests heavily invaded by *A. dealbata*, in central Portugal, to test the hypotheses that riparian forest invasion would negatively affect macroinvertebrate abundance, taxa richness (total and of sensitive taxa), and diversity, as well as change community taxonomic structure. Community structure in terms of functional feeding groups was also hypothesized to differ between stream types, especially for shredders that strongly depend on allochthonous leaf litter and have their life cycles synchronized with the autumnal leaf litterfall in native forests. Since *A. dealbata* is widely naturalized in all southern Europe (Sheppard et al., 2006; Lorenzo et al., 2010), this study contributes with relevant information that can inform about the cross-ecosystem effects of riparian invasion on stream communities in Mediterranean countries.

METHODS

Study area and streams

This study took place in Serra da Lousã, central Portugal, where invasive *Acacia* species (mainly



Figure 1. Image of a valley (São João stream) in the study area in Serra da Lousã, central Portugal, in February 2019, depicting heavy invasion by *A. dealbata*. *A. dealbata* is easily recognizable by its bright yellow flowers. *Imagen de un valle (arroyo São João) en el área de estudio en la Serra da Lousã, centro de Portugal, en febrero del 2019, que muestra una fuerte invasión de A. dealbata. A. dealbata es fácilmente reconocible por sus flores de color amarillo brillante.*

Acacia dealbata Link.) cover large areas, replacing native shrub communities and native deciduous forests dominated by *Quercus* spp. and *Castanea sativa* Mill. (Ferreira *et al.*, 2021; Pereira *et al.*, 2021). Six small streams (< 4 m wide, basin area ≤ 6 km²) were selected: three streams flow through native mixed deciduous forests (native streams: Maior, Cerdeira and Candal) and the other three streams flow through forests heavily invaded by *A. dealbata* (cover in the riparian area: 94 – 100%) (invaded streams: Sotão, Fiscal and Piedade) (Table 1). All streams have low conductivity and circumneutral pH, and the substrate is composed mainly of schist gravel and cobbles, with coarse sand in depositional areas (Pereira & Ferreira, 2021).

Stream water was characterized regarding dissolved nutrient concentrations on three dates (3 January, 14 January and 11 February 2019) preceding the macroinvertebrate sampling (26 February 2019). Stream water was collected, filtered in the field through fiberglass filters (47 mm diameter, 0.7 μ m pore size; Whatman GF/F, GE Healthcare UK Limited, Little Chalfont, UK) and

distributed by two acid-washed flasks for determination of soluble reactive phosphorus (SRP) concentration (ascorbic acid method; APHA, 1995) and of nitrate, nitrite and ammonia concentrations (colorimetric method; AA3 Bran + Luebbe autoanalyzer; SEAL Analytical, Norderstedt, Germany). Dissolved inorganic nitrogen (DIN) concentrations were estimated as the sum of the concentrations of nitrate-N, nitrite-N and ammonia-N.

Benthic macroinvertebrate communities

Benthic macroinvertebrates were sampled from all streams on 26 February 2019 using a semi-quantitative method. Three samples were taken over a representative 50-m section in each stream by disturbing (kicking) the substrate and allowing the current to move the dislodged macroinvertebrates into a hand net (30 \times 30 cm opening, 0.5 mm mesh size) that was placed just downstream of the operator (the same across streams); kicking was done from downstream to upstream for a standard distance and time (1 m for 1 minute)

Table 1. Location, basin area, *Acacia dealbata* cover in the riparian area, and water nutrient concentrations (mean \pm SD, $n = 3$ between 3 January and 11 February 2019) of streams flowing through native forests (native streams) and of streams flowing through forests heavily invaded by *A. dealbata* (invaded streams) in Serra da Lousã, central Portugal. Comparison between stream types was done by t-test considering $\alpha = 0.050$, and p -values are shown (significant values are highlighted in bold). *Localización, área de la cuenca, A. dealbata en la zona de ribera, y concentraciones de nutrientes en el agua (media \pm DE, $n = 3$ entre el 3 de enero y el 11 de febrero del 2019) de arroyos que fluyen a través de bosques nativos (arroyos nativos) y de arroyos que fluyen a través de bosques fuertemente invadidos por A. dealbata (arroyos invadidos) en la Sierra de Lousã, centro de Portugal. La comparación entre tipos de arroyos se realizó mediante t-test considerando $\alpha = 0.050$, y se muestran los p-valores (los valores significativos se resaltan en negrita).*

Stream	Latitude (N)	Longitude (W)	Altitude (m a.s.l)	Basin area (km ²) ¹	Riparian <i>Acacia dealbata</i> (%) ^{1,2}	DIN (μ g N/L) ³	SRP (μ g P/L) ⁴
<i>Native streams</i>							
Maior	40°07'53.3"	8°11'40.7"	195	6.05	1	25 \pm 4	9 \pm 6
Cerdeira	40°05'23.1"	8°12'05.0"	529	3.37	1	22 \pm 2	8 \pm 5
Candal	40°04'54.1"	8°12'16.6"	634	0.98	0	20 \pm 2	7 \pm 2
<i>Invaded streams</i>							
Sotão	40°07'54.1"	8°09'08.3"	373	1.14	94	76 \pm 28	23 \pm 7
Fiscal	40°06'40.2"	8°13'35.1"	329	1.24	100	60 \pm 17	14 \pm 2
Piedade	40°05'52.6"	8°14'11.5"	250	2.15	100	47 \pm 8	9 \pm 5
<i>Average</i>							
Native streams			453	3.47	1	23 \pm 3	8 \pm 1
Invaded streams			317	1.51	98	61 \pm 15	15 \pm 7
t-test (p-value)			0.380	0.262	<0.001	0.012	0.139

¹ from Pereira *et al.* (2021); ² the riparian area was defined as an area 50 m wide on each stream bank and 250 m long upstream from the sampling point;

³ DIN, dissolved inorganic nitrogen; ⁴ SRP, soluble reactive phosphorus

and samples were collected moving from downstream to upstream in the stream section. Benthic samples were placed in plastic bags with stream water and transported to the laboratory in cold boxes. In the laboratory, samples were preserved with formaldehyde (final concentration of 4%) and stored in the dark until processed. Samples were rinsed thoroughly with tap water on top of a series of sieves (2 cm, 1 cm, and 0.5 mm mesh) to fractionate the sample and facilitate subsequent macroinvertebrate sorting. After sorting, macroinvertebrates were kept in vials with 80% ethanol. Macroinvertebrates were then identified at the lowest taxonomic level possible (Tachet et al., 2000; Vieira-Lanero, 2000) and counted under a stereo microscope (Leica, M80, Wetzlar, Germany). Macroinvertebrates were also grouped by functional feeding groups (FFG): predators, shredders, grazers/scrapers, gatherers/collectors, and filter feeders (Tachet et al., 2000).

Data analysis

Altitude, basin area, and water nutrient concentrations (SRP and DIN; average of the three dates) were compared between stream types (native vs. invaded, $n = 3$) by student's t-test ($\alpha = 0.050$). Macroinvertebrate abundance, total taxa richness, Ephemeroptera + Plecoptera + Trichoptera (EPT, known as sensitive taxa) taxa richness, diversity indices, and abundance of FFG (average of the three samples) were also compared between stream types (native vs. invaded, $n = 3$) by student's t-test ($\alpha = 0.050$). Two diversity indices were used: the Margalef species richness index (d), which is sensitive to abundant species (Morris et al., 2014), has its maximum value when all individuals belong to different species and its minimum when all individuals belong to the same species, i.e., a higher value indicates higher diversity; and the Pielou species evenness index, where a higher value indicates higher uniformity in the distribution of individuals by species and therefore higher diversity (Pielou, 1996).

Macroinvertebrate communities (taxonomic and FFG matrices) were compared between stream types (native vs. invaded) by analysis of similarity (ANOSIM, $\alpha = 0.050$) based on a Bray-Curtis similarity matrix of log-transformed

abundance data; visual representation of the similarity among communities was achieved by non-metric multidimensional scaling (nMDS; considered valid if stress level < 0.20). The taxa or FFGs responsible for distinguishing between stream types were identified by performing similarity percentage (SIMPER) analysis, which examined the contribution of each taxon or FFG to the mean dissimilarity between stream types.

Student's t-tests were performed using Excel (Microsoft Office 2019) and diversity indices and multivariate analyses were performed using PRIMER 6 (v6.1.16) & PERMANOVA+ (v1.0.6; Primer-E Ltd, Plymouth, UK) software.

RESULTS

Study streams

Mean dissolved nutrient concentrations were $1.9\times$ (SRP) and $2.7\times$ (DIN) higher in invaded than in native streams, but the difference was significant only for DIN (Table 1). Still, DIN concentrations were in the oligotrophic range in all streams ($< 77 \mu\text{g/L}$) (Table 1).

Benthic macroinvertebrate communities

In total, benthic samples retrieved 8599 macroinvertebrates distributed by 84 taxa (Table S1, see Supplementary information at <https://www.limnetica.net/en/limnetica>). Mayflies (Ephemeroptera: 32% individuals, 11 taxa), caddisflies (Trichoptera: 17% individuals, 22 taxa), worms (Oligochaeta: 16% individuals, 5 taxa), true flies (Diptera: 13% individuals, 16 taxa), and stoneflies (Plecoptera: 12% individuals, 8 taxa) were the most represented groups across streams (Table S1).

Mean abundance was $2.5\times$ higher in native than in invaded streams, but differences were non-significant due to high variation (up to $5.3\times$) among native streams (Table 2). A similar pattern was observed for total taxa richness, which was 20% higher in native than in invaded streams, with differences being non-significant due to high variation (25%) among native streams (Table 2). However, stream types significantly differed in EPT taxa richness, which was 47% higher in

Table 2. Benthic macroinvertebrate community parameters (mean \pm SE, $n = 3$) in streams flowing through native forests (native streams) and in streams flowing through forests heavily invaded by *A. dealbata* (invaded streams) in Serra da Lousã, central Portugal, in February 2019. Comparison between stream types was done by t-test considering $\alpha = 0.050$, and p -values are shown (significant values are highlighted in bold). *Parámetros de la comunidad de macroinvertebrados bentónicos (media \pm DE, $n = 3$) en arroyos que fluyen a través de bosques nativos (arroyos nativos) y en arroyos que fluyen a través de bosques fuertemente invadidos por *A. dealbata* (arroyos invadidos) en la Sierra de Lousã, centro de Portugal, en febrero del 2019. La comparación entre tipos de arroyos se realizó mediante t-test considerando $\alpha = 0.050$, y se muestran los p-valores (los valores significativos se resaltan en negrita).*

Stream	Abundance (no. ind./sample)	Total taxa richness (no. taxa/sample)	EPT taxa richness (no. taxa/sample)	Margalef index (d)	Pielou index (J')
<i>Native streams</i>					
Maior	728 \pm 134	40 \pm 1	23 \pm 1	5.91 \pm 0.26	0.74 \pm 0.00
Cerdeira	1115 \pm 150	40 \pm 5	22 \pm 1	5.63 \pm 0.73	0.68 \pm 0.09
Candal	210 \pm 77	32 \pm 4	20 \pm 2	5.92 \pm 0.41	0.83 \pm 0.04
<i>Invaded streams</i>					
Sotão	366 \pm 27	35 \pm 1	14 \pm 2	5.71 \pm 0.26	0.75 \pm 0.02
Fiscal	238 \pm 28	30 \pm 0	15 \pm 2	5.32 \pm 0.12	0.76 \pm 0.02
Piedade	209 \pm 82	30 \pm 5	17 \pm 3	5.56 \pm 0.49	0.81 \pm 0.05
<i>Average</i>					
Native streams	684 \pm 22	37 \pm 1	22 \pm 1	5.82 \pm 0.14	0.75 \pm 0.02
Invaded streams	271 \pm 18	31 \pm 1	15 \pm 1	5.53 \pm 0.11	0.77 \pm 0.01
t-test (p -value)	0.196	0.132	0.010	0.121	0.694

EPT, Ephemeroptera + Plecoptera + Trichoptera (see Table S1)

native than invaded streams (Table 2). Diversity indices did not significantly differ between stream types (Table 2).

Macroinvertebrate community structure (based on the taxonomic matrix) significantly differed between native and invaded streams (ANOSIM, $R = 0.47$, $p = 0.001$) (Fig. 2). The dissimilarity of macroinvertebrate communities between stream types was moderate (SIMPER analysis, cumulative dissimilarity = 61%). The taxa that contributed the most to differences between native and invaded streams (contribution to dissimilarity $> 2.5\%$) were *Amphinemura* sp., *Habroleptoides* sp., *Chloroperla* sp., and *Rithrogena* sp., which were more abundant in native streams, and *Ephemerella* sp. Simuliidae, and *Paraleptophlebia* sp., which were more abundant in invaded streams (Table 3).

Abundance of FFGs (predators, shredders, grazers/scrapers, gatherers/collectors, and filter feeders) did not significantly differ between native and invaded streams (t-tests, $p \geq 0.111$; Fig. 3A). Still, macroinvertebrate communities (based on the FFG matrix) significantly differed between native and invaded streams (ANOSIM, $R = 0.25$, $p = 0.012$), owing to the slightly higher relative

contribution of shredders and scrapers/grazers in native streams and of filter feeders in invaded streams (Fig. 3B), but dissimilarity of macroinvertebrate communities between stream types was low (SIMPER analysis, cumulative dissimilarity = 16%).




Figure 2. Ordination (nMDS based on Bray-Curtis similarity of log-transformed abundance data) of benthic macroinvertebrate communities in streams flowing through native forests (native streams) and in streams flowing through forests heavily invaded by *A. dealbata* (invaded streams) in Serra da Lousã, central Portugal, in February 2019. *Ordenación (nMDS basada en la similitud Bray-Curtis de datos de abundancia transformados por logaritmo) de las comunidades de macroinvertebrados bentónicos en arroyos que fluyen por bosques nativos (arroyos nativos) y en arroyos que fluyen por bosques fuertemente invadidos por *A. dealbata* (arroyos invadidos) en la Sierra de Lousã, centro de Portugal, en febrero del 2019.*

Table 3. Result of the similarity percentage (SIMPER) analysis, performed on benthic macroinvertebrate communities, showing the contribution (%) of each taxon to the dissimilarity between streams flowing through native forests (native streams) and streams flowing through forests heavily invaded by *A. dealbata* (invaded streams) in Serra da Lousã, central Portugal, in February 2019. *Resultado del análisis de porcentaje de similitud (SIMPER), realizado en comunidades de macroinvertebrados bentónicos, mostrando la contribución (%) de cada taxon a la disimilitud entre arroyos que fluyen por bosques nativos (arroyos nativos) y arroyos que fluyen por bosques fuertemente invadidos por A. dealbata (arroyos invadidos) en la Sierra de Lousã, centro de Portugal, en febrero del 2019.*

Taxa	Native average abundance	Invaded average abundance	Average dissimilarity	Dissimilarity/SD	Contribution %	Cumulative %
<i>Amphinemura</i> sp.	3.49	0.20	2.60	3.32	4.29	4.29
<i>Habroleptoides</i> sp.	3.20	0.00	2.32	1.35	3.84	8.12
<i>Chloroperla</i> sp.	2.76	0.00	2.13	2.75	3.51	11.63
<i>Ephemera</i> sp.	0.15	2.60	2.03	3.09	3.36	14.99
Simuliidae	0.75	2.40	1.64	1.37	2.72	17.71
<i>Rithrogenea</i> sp.	2.42	0.63	1.64	1.50	2.71	20.41
<i>Paraleptophlebia</i> sp.	1.26	1.60	1.58	1.12	2.61	23.03
<i>Ecdyonurus</i> sp.	2.94	1.53	1.52	1.66	2.50	25.53
<i>Acentrella sinaica</i>	2.85	1.96	1.39	1.33	2.30	27.83
<i>Lepidostoma hirtum</i>	2.13	0.88	1.38	1.45	2.28	30.11
<i>Larcasia partita</i>	2.15	1.02	1.37	1.49	2.26	32.37
<i>Baetis</i> sp.	2.25	0.99	1.33	1.45	2.20	34.58
<i>Polycelis</i> spp.	1.49	1.36	1.30	1.19	2.15	36.73
<i>Epeorus</i> sp.	1.77	0.77	1.30	1.27	2.14	38.87
<i>Duguesia</i> spp.	1.50	0.80	1.29	1.12	2.13	41.00
<i>Beraea</i> sp.	1.82	0.38	1.29	1.19	2.13	43.13
<i>Hydrocyphon</i> sp.	1.23	1.14	1.19	1.33	1.96	45.09
<i>Heptagenia</i> sp.	1.55	0.18	1.18	1.52	1.95	47.04
Lumbriculidae	2.73	3.17	1.16	1.24	1.91	48.96
<i>Diplectrona felix</i>	1.23	0.92	1.05	1.08	1.74	50.69
<i>Hydropsyche</i> sp.	2.41	2.30	1.02	1.28	1.68	52.37
<i>Protonemura</i> sp.	0.86	1.45	0.91	1.29	1.50	53.87
<i>Thremma</i> sp.	1.50	0.48	0.90	1.46	1.49	55.36
Tanytarsini	2.61	2.28	0.90	1.58	1.49	56.85
<i>Sphaerium</i> sp.	0.20	1.19	0.88	1.45	1.45	58.30
Orthocladinae	1.89	1.60	0.85	1.48	1.41	59.71
<i>Leuctra</i> sp.	0.76	0.91	0.85	1.15	1.40	61.11

DISCUSSION

In this study we assessed the effects of the invasion of native mixed deciduous forests by *A. dealbata* on stream benthic macroinvertebrate communities in central Portugal. Following the pattern found in the literature (see examples in the Introduction), invasion effects were more evident for macroinvertebrate community structure than for abundance, total richness or diversity indices.

In accordance with our hypothesis, macroinvertebrate community structure differed between stream types, which agrees with other studies addressing invasion effects on benthic macroinvertebrate communities (Lowe et al., 2008; Serra et al., 2013; Seeney et al., 2019; Little et al., 2021). In particular, we found higher EPT taxa richness and higher number of EPT taxa as indicators for streams flowing through native than invaded forests, as also seen in the case of *L. maacki* (Little et




Figure 3. Benthic macroinvertebrates functional feeding groups (FFG) abundance (A) and relative contribution (B) in streams flowing through native forests (native streams) and in streams flowing through forests heavily invaded by *A. dealbata* (invaded streams) in Serra da Lousã, central Portugal, in February 2019. *Abundancia (A) y contribución relativa (B) de grupos funcionales de alimentación (FFG) de macroinvertebrados bentónicos en arroyos que fluyen por bosques nativos (arroyos nativos) y en arroyos que fluyen por bosques fuertemente invadidos por A. dealbata (arroyos invadidos) en la Sierra de Lousã, centro de Portugal, en febrero del 2019.*

al., 2021) and *A. mearnsii* (Magoba & Samways, 2010) invasions. EPT orders are known to have many taxa that are sensitive to environmental perturbations (Fore, 1996). Concurrently, invaded sites had higher contribution of generalist taxa, such as Simuliidae and Sphaeridae (Fore, 1996; Tampo *et al.*, 2021).

Macroinvertebrate community structure differed between stream types also when macroinvertebrate FFGs were considered. Shredders strongly depend on allochthonous leaf litter as a food resource (Wallace *et al.*, 1997). In this sense, vegetation changes can produce important variations in the distribution of FFG (Balderas *et al.*, 2016). As seen in Mancilla *et al.* (2009), in central

Chile, streams flowing through basins with less than 20% native vegetation cover had significantly higher number of gatherers and lower number of shredders than streams flowing through basins with higher percentage of native vegetation. Shredders transform coarse organic resources into fine particulate organic matter, thus supporting secondary consumers (Cummins *et al.*, 1973; Patrick, 2013), such as gatherers/collectors and filter feeders (Shepard & Minshall, 1984). Alterations in forest composition that result in decreased quality of leaf litter inputs to streams and augmented canopy cover can lead to decreases in litter decomposition rates and algal production, diminishing resources availability and inducing shifts in consumer assemblages, as found in the case of invasion of deciduous riparian forests by *Rhododendron ponticum* L. in Western Ireland (Hladyz *et al.*, 2011). Similar changes resulting from the invasion of deciduous forests by *A. dealbata* can explain the tendency for the lower abundance of grazers/scrapers and higher abundance of filter feeders in invaded streams.

Contrary to anticipated, no significant differences were found in macroinvertebrate abundance, total taxa richness, and diversity between streams flowing through native forests and streams flowing through forests heavily invaded by *A. dealbata*, which was mainly attributed to high variation among native streams. Native streams may have differed more in current velocity or in the composition of plant litter input than invaded streams that had their riparian vegetation dominated by *A. dealbata* (Pereira *et al.*, 2021).

Our data suggests that invasion of native deciduous forests by *A. dealbata* may affect community taxonomic and functional (FFG) structure, although it may not strongly affect macroinvertebrate communities in terms of total abundance, total richness, and diversity. However, we recall that benthic macroinvertebrate sampling was performed only once, in winter. Benthic macroinvertebrate communities can exhibit natural seasonal variation, induced by various factors such as food availability, temperature, and photoperiod, which may moderate invasion effects (Álvarez-Cabria *et al.*, 2011; Seeney *et al.*, 2019). Additionally, the use of a unique sampling time did not allow assessing successional changes occurring over time

during the invasion process, shading temporal dynamics and the impact of different percentages of *Acacia* cover. Moreover, lack of pre-invasion data makes it challenging to establish a cause-and-effect relationship between the invasion and the observed changes in community structure. Finally, the reduced number of streams used may have contributed to the high variability within stream types (especially among native streams), precluding the identification of clearer invasion effects.

Future studies should consider a larger number of streams to take into consideration the natural variability among native streams and the different invasion degrees in invaded streams; this could allow establishing correlations between the degree of invasion and macroinvertebrate metrics (Ferreira et al., 2016). Also, benthic macroinvertebrate sampling should be carried out seasonally to take into consideration macroinvertebrate life cycles and litter inputs phenology to streams, which may moderate invasion effects on benthic macroinvertebrates (Seeney et al., 2019). Additionally, since invasion by *A. dealbata* is proceeding at high rates (500 ha/year between 2005 and 2015, and likely faster since 2017 owing to large-scale wild fires; ICNF, 2019; Ferreira et al., 2021), pre-invasion information may (soon) become available for streams that integrate the official bioassessment scheme and that were sampled before invasion (early 2000's) and that now flow through forests which became invaded; this would allow before-after comparisons. Finally, since *A. dealbata* is invasive in Mediterranean countries, collaborative assessments should also be considered for a large scale understanding of invasion impacts on stream communities. A better understanding of invasion impacts on streams could help identify most vulnerable ecosystems and envisage mitigation measures.

ACKNOWLEDGEMENTS

This study was financed by the Portuguese Foundation for Science and Technology (FCT) through the strategic projects UIDP/04292/2020 and UIDB/04292/2020 granted to MARE and LA/P/0069/2020 granted to the Associate Laboratory ARNET, and through financial support given to VF (IF/00129/2014, CEECIND/02484/2018).

SGA undertook this study in the scope of the course “Projecto em Biologia Aplicada II” (undergraduate studies in Biology, Department of Life Sciences, University of Coimbra). Ana Pereira is thanked for stream water collection. Water nutrient analyses were ordered to the Centro de Apoio Científico-Tecnológico á Investigación (CACTI, University of Vigo, Spain).

REFERENCES

- Abelho, M. & Graça, M. A. S. (1996). Effects of eucalyptus afforestation on leaf litter dynamics and macroinvertebrate community structure of streams in central Portugal. *Hydrobiologia*, 324, 195–204. DOI:10.1007/BF00016391
- Álvarez-Cabria, M., Barquín, J. & Juanes, J. A. (2011). Macroinvertebrate community dynamics in a temperate European Atlantic river. Do they conform to general ecological theory? *Hydrobiologia*, 658(1), 277–291. DOI: 10.1007/s10750-010-0498-y
- Atwood, T. B., Wiegner, T. N., Turner, P. & MacKenzie, R. A. (2010). Potential effects of an invasive nitrogen-fixing tree on a Hawaiian stream food web. *Pacific Science*, 64(3), 367–379. DOI: 10.1215/23112
- Balderas, E. C. S., Grac, C., Berti-Equille, L. & Hernandez, M. (2016). Potential application of macroinvertebrates indices in bioassessment of Mexican streams. *Ecological Indicators*, 61, 558–567. DOI: 10.1016/j.ecolind.2015.10.007
- Beschta, R. L. (1997). Riparian shade and stream temperature: An alternative perspective. *Rangelands*, 19, 25–28. DOI: 10.1015/640452
- Castro-Díez, P., Godoy, O., Alonso, A., Gallardo, A. & Saldaña, A. (2014). What explains variation in the impacts of exotic plant invasions on the nitrogen cycle? A meta-analysis. *Ecology Letters*, 17(1), 1–12. DOI: 10.1111/ele.12197
- Cummins, K.W. (1973). Trophic relations of aquatic insects. *Annual Review of Entomology*, 18, 183–206. DOI: 10.1146/annurev.en.18.010173.001151
- Dosskey, M. G., Vidon, P., Gurwick, N. P., Allan, C. J., Duval, T. P. & Lowrance, R. (2010). The role of riparian vegetation in protecting and improving chemical water quality in streams.

- Journal of the American Water Resources Association*, 46, 261–277. DOI: 10.1111/j.1752-1688.2010.00419.x
- Ferreira, V., Castela, J., Rosa, P., Tonin, A. M., Boyero, L. & Graça, M. A. S. (2016). Aquatic hyphomycetes, benthic macroinvertebrates and leaf litter decomposition in streams naturally differing in riparian vegetation. *Aquatic Ecology*, 50(4), 711–725. DOI: 10.1007/s10452-016-9588-x
- Ferreira, V., Figueiredo, A., Graça, M. A. S., Marchante, E. & Pereira, A. (2021). Invasion of temperate deciduous broadleaf forests by N-fixing tree species – Consequences for stream ecosystems. *Biological Reviews*, 96, 877–902. DOI: 10.1111/brv.12682
- Fore, L. S., Karr, J. R. & Wisseman, R. W. (1996). Assessing invertebrate responses to human activities: Evaluating alternative approaches. *Journal of the North American Benthological Society*, 15(2), 212–231. DOI: 10.2307/1467949
- Gallagher, R. V., Leishman, M. R., Miller, J. T., Hui, C., Richardson, D. M., Suda, J. & Trávníček, P. (2011). Invasiveness in introduced Australian acacias: The role of species traits and genome size. *Diversity and Distributions*, 17, 884–897. DOI: 10.1111/j.1472-4642.2011.00805.x
- Gerhard, M. & Reich, M. (2000). Restoration of streams with large wood: Effects of accumulated and built-in wood on channel morphology, habitat diversity and aquatic fauna. *International Review of Hydrobiology*, 85(1), 123–137. DOI:10.1002/(SICI)1522-2632(200003)85:1<123::AID-IROH123>3.0.CO;2-T
- Goldstein, C. L., Williard, K. W. & Schoonover, J. E. (2009). Impact of an invasive exotic species on stream nitrogen levels in Southern Illinois. *Journal of the American Water Resources Association*, 45(3), 664–672. DOI: 10.1111/j.1752-1688.2009.00314.x
- Hawkins, C. P., Murphy, M. L. & Anderson, N. H. (1982). Effects of canopy, substrate composition, and gradient on the structure of macroinvertebrate communities in Cascade Range streams of Oregon. *Ecology*, 63, 1840–1856. DOI: 10.2307/1940125
- Hladyz, S., Åbjörnsson, K., Giller, P. S. & Woodward, G. (2011). Impacts of an aggressive riparian invader on community structure and ecosystem functioning in stream food webs. *Journal of Applied Ecology*, 48(2), 443–452. DOI: 10.1111/j.1365-2664.2010.01924.x
- ICNF (2019) *IFN6 – Principais resultados – relatório sumário*. Instituto da Conservação da Natureza e das Florestas. Lisboa, Portugal. <http://www.icnf.pt/portal/florestas/ifn/ifn5/>
- Kominoski, J. S., Shah, J. J. F., Canhoto, C., Fischer, D. G., Giling, D. P., González, E., ... Tiegs, S. D. (2013). Forecasting functional implications of global changes in riparian plant communities. *Frontiers in Ecology and the Environment*, 11(8), 423–432. DOI: 0.1890/120056
- Lecerf, A., Patfield, D., Boiché, A., Riipinen, M. P., Chauvet, E. & Dobson, M. (2007). Stream ecosystems respond to riparian invasion by Japanese knotweed (*Fallopia japonica*). *Canadian Journal of Fisheries and Aquatic Sciences*, 64(9), 1273–1283. DOI: 10.1139/f07-092
- Little, M., Custer, K. W., Borth, E. B., Chapman, J. I., Kukla, M. J., Kuminecz, C., ... McEwan, R. W. (2021). The influence of riparian invasion by the terrestrial shrub *Lonicera maackii* on aquatic macroinvertebrates in temperate forest headwater streams. *Biological Invasions*, 23(1), 25–35. DOI: 10.1007/s10530-020-02349-8
- Lorenzo, P., González, L. & Reigosa, M. J. (2010). The genus *Acacia* as invader: The characteristic case of *Acacia dealbata* Link in Europe. *Annals of Forest Science*, 67, 101. DOI: 10.1051/forest/2009082
- Lorenzo, P., Pazos-Malvido, E., González, L. & Reigosa, M. J. (2008). Allelopathic interference of invasive *Acacia dealbata*: Physiological effects. *Allelopathy Journal*, 22, 452–462. DOI:10.1007/s11258-010-9831-9
- Lowe, S., Woodford, D. J., Impson, D. & Day, J. A. (2008). The impact of invasive fish and invasive riparian plants on the invertebrate fauna of the Rondegat River, Cape Floristic Region, South Africa. *African Journal of Aquatic Science*, 33(1), 51–62. DOI: 10.2989/ajas.2007.33.1.6.390

- Magoba, R. N. & Samways, M. J. (2010). Recovery of benthic macroinvertebrate and adult dragonfly assemblages in response to large scale removal of riparian invasive alien trees. *Journal of Insect Conservation*, 14(6), 627–636. DOI: 10.1007/s10841-010-9291-5
- Mancilla, G., Valdovinos, C., Azócar, M., Jorquera, P. & Figueroa, R. (2009). Efecto del reemplazo de la vegetación nativa de ribera sobre la comunidad de macroinvertebrados bentónicos en arroyos de climas templados, Chile central. *Hidrobiológica*, 19(3), 193–203.
- Masese, F. O., Kitaka, N., Kipkemboi, J., Gettel, G. M., Irvine, K. & McClain, M. E. (2014). Macroinvertebrate functional feeding groups in Kenyan highland streams: Evidence for a diverse shredder guild. *Freshwater Science*, 33(2), 435–450. DOI: 10.1086/675681
- Marchante, E., Kjølner, A., Struwe, S. & Freitas, H. (2008). Short- and long-term impacts of *Acacia longifolia* invasion on the below-ground processes of a Mediterranean coastal dune ecosystem. *Applied Soil Ecology*, 40(2), 210–217. DOI: 10.1016/j.apsoil.2008.04.004
- Mineau, M. M., Baxter, C. V., Marcarelli, A. M. & Minshall, G. W. (2012). An invasive riparian tree reduces stream ecosystem efficiency via a recalcitrant organic matter subsidy. *Ecology*, 93(7), 1501–1508. DOI: 10.1890/11-1700.1
- Molinero, J. & Pozo, J. (2004). Impact of a eucalyptus (*Eucalyptus globulus* Labill.) plantation on the nutrient content and dynamics of coarse particulate organic matter (CPOM) in a small stream. *Hydrobiologia*, 528(1–3), 143–165. DOI: 10.1007/s10750-004-2338-4
- Morris, E. K., Caruso, T., Buscot, F., Fisher, M., Hancock, C., Maier, T. S., ... Rillin, M. C. (2014). Choosing and using diversity indices: Insights for ecological applications from the German Biodiversity Exploratories. *Ecology and Evolution*, 4(18), 3514–3524. DOI: 10.1002/ece3.1155
- Patrick, C. J. (2013). The effect of shredder community composition on the production and quality of fine particulate organic matter. *Freshwater Science*, 32(3), 1026–1035. DOI: 10.1899/12-090.1
- Pereira, A. & Ferreira, V. (2021). Invasion of native riparian forests by *Acacia* species affects in-stream litter decomposition and associated microbial decomposers. *Microbial Ecology*, 81, 14–25. DOI: 10.1007/s00248-020-01552-3
- Petsch, D. K. (2016). Causes and consequences of biotic homogenization in freshwater ecosystems. *International Review of Hydrobiology*, 101(3–4), 113–122. DOI: 10.1002/iroh.201601850
- Pielou, E. C. (1966). The measurement of diversity in different types of biological collections. *Journal of Theoretical Biology*, 13, 131–144. DOI: 10.1016/0022-5193(66)90013-0
- Pohlman, C. L., Nicotra, A. B. & Murray, B. R. (2005). Geographic range size, seedling ecophysiology and phenotypic plasticity in Australian *Acacia* species. *Journal of Biogeography*, 32, 341–351. DOI: 10.1111/j.1365-2699.2004.01181.x
- Seeney, A., Pattison, Z., Willby, N. J., Boon, P. J. & Bull, C. D. (2019). Stream invertebrate diversity reduces with invasion of river banks by non-native plants. *Freshwater Biology*, 64(3), 485–496. DOI: 10.1111/fwb.13236
- Serra, M. N., Albariño, R. & Díaz Villanueva, V. (2013). Invasive *Salix fragilis* alters benthic invertebrate communities and litter decomposition in northern Patagonian streams. *Hydrobiologia*, 701, 173–188. DOI: 10.1007/s10750-012-1270-2
- Shepard, R. K. & Minshall, G. W. (1984). Role of benthic insect feces in a Rocky Mountain stream: Fecal production and support of consumer growth. *Ecography*, 7(2), 119–127. DOI: 10.1111/j.1600-0587.1984.tb01112.x
- Sheppard, A. W., Shaw, R. H. & Sforza, R. (2006). Top 20 environmental weeds for classical biological control in Europe: A review of opportunities, regulations and other barriers to adoption. *Weed Research*, 46, 93–117. DOI: 10.1111/j.1365-3180.2006.00497.x
- Stewart, S. D., Young, M. B., Harding, J. S. & Horton, T. W. (2019). Invasive nitrogen-fixing plant amplifies terrestrial–aquatic nutrient flow and alters ecosystem function. *Ecosystems*, 22, 587–601. DOI: 10.1007/s10021-018-0289-2
- Tachet, H., Richoux, P., Bournand, M. & Usseglio-Polatera, P. (2000). *Invertébrés d'eau*

- douce: Systématique, biologie, écologie*. CNRS Editions. France.
- Tampo, L., Kaboré, I., Alhassan, E. H., Ouéda, A., Bawa, L. M. & Djaneye-Boundjou, G. (2021). Benthic macroinvertebrates as ecological indicators: Their sensitivity to the water quality and human disturbances in a tropical river. *Frontiers in Water*, 3. DOI: 10.3389/frwa.2021.662765
- Tolkkinen, M. J., Heino, J., Ahonen, S. H. K., Lehosmaa, K. & Mykrä, H. (2020). Streams and riparian forests depend on each other: A review with a special focus on microbes. *Forest Ecology and Management*, 462, 117962. DOI: 10.1016/j.foreco.2020.117962
- Vieira-Lanero, R. (2000). *Las larvas de los tricópteros de Galicia (Insecta: Trichoptera)*. (PhD. Thesis. Universidad de Santiago de Compostela, España). Retrieved: <https://dialnet.unirioja.es/servlet/tesis?codigo=71023>
- Wallace, J. K., Eggert, S. L., Meyer, J. & Webster, J. S. (1997). Multiple trophic levels of a forest stream linked to terrestrial litter inputs. *Science*, 277(5322), 102–104. DOI: 10.1126/science.277.5322.102